

**Escherichia Coli  
Antibody**  
Equine Origin  
**Bar-Guard-99™**  
For Oral Use Only  
1 Dose/10mL


**Directions:** Shake well and use immediately. See enclosure for complete directions and precautions.

**Dosage:** Using aseptic technique, administer 10 mL (calves) or 2 mL (lambs) orally as soon as possible after birth. For optimal results, Bar-Guard-99™ should be administered after colostrum but within 12 hours after birth and preferably within 6 hours of birth. Bar-Guard-99™ may also be administered within 8 hours of exposure, not to exceed 20 hours of age. Contains cresol as a preservative.


**Precautions:** Store in dark at 35-45°F. Do not administer within 21 days before slaughter. Anaphylactoid reactions may occur.

**Antidote:** Epinephrine.

Boehringer Ingelheim Vetmedica, Inc.  
St. Joseph, MO 64506 USA  
U.S. Vet Lic. No. 124  
408109N-00



Ser. No.: \_\_\_\_\_  
Exp. Date: \_\_\_\_\_



0 12313 10811 3

Serial No.: \_\_\_\_\_  
Exp. Date: \_\_\_\_\_


**Escherichia Coli  
Antibody**

Equine Origin  
**Bar-Guard-99™**  
For Oral Use Only  
5 Dose/50 mL

**Lambs:** Using aseptic technique, administer 2 mL orally as soon as possible after birth. Bar-Guard-99 should be administered within 12 hours after birth and preferably within 6 hours of birth. Contains cresol as a preservative. Anaphylactoid reactions may occur.

**Antidote:** Epinephrine.

**Boehringer Ingelheim  
Vetmedica, Inc.**  
St. Joseph, MO 64506 U.S.A.  
U.S. Veterinary License No. 124  
Code 108-121; 08102-02



**Boehringer Ingelheim Vetmedica, Inc.**  
2621 North Belt Highway • St. Joseph, MO 64506-2002  
Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717  
[www.bi-vetmedica.com](http://www.bi-vetmedica.com)