

Exp. Date:  
Serial No.:

**Indications:** For the vaccination of all types and ages of healthy horses against Eastern, Western and Venezuelan Equine Encephalomyelitis and tetanus.

**Administration and Dosage:** Shake well before using. Inject 2 mL deep into muscle. Revaccination required. See carton for full directions.

**Caution:** Store at 35-45°F (2-7°C). Protect from freezing. Use entire contents when first opened. Contains gentamicin and thimerosal as

## Encephalomyelitis Vaccine,

Eastern, Western, and Venezuelan, Killed Virus

## Tetanus Toxoid

**Cephalovac® VEWT**

10 Doses/20 mL

preservatives. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur following use.

**Antidote:** Epinephrine.

**Boehringer Ingelheim Vetmedica, Inc.**

**St. Joseph, MO 64506 U.S.A.**

**U.S. Veterinary License No. 124**

Code 131-325; 31303-02



**Boehringer  
Ingelheim**

**Boehringer Ingelheim Vetmedica, Inc.**

**2621 North Belt Highway • St. Joseph, MO 64506-2002**

**Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717**

**[www.bi-vetmedica.com](http://www.bi-vetmedica.com)**