

Exp. Date:
Serial No.:

Indications: For vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the prevention of lymphoid depletion, inflammation and colonization of lymphoid tissue associated with Porcine Circovirus Type 2 (PCV2). The onset of protection is 2 weeks post-vaccination. The duration of immunity is at least 17 weeks.

Directions and Dosage: See carton.

Precautions: Store at 35-45°F (2-7°C). Do not freeze. Shake well before using. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactic reaction, epinephrine is symptomatic treatment.

Preservative: Neomycin.



Porcine Circovirus Vaccine

Type 2, Killed Baculovirus Vector

Ingelvac CircoFLEX®

250 Doses/250 mL

Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, MO 64506
U.S. Veterinary License No. 124



Boehringer
Ingelheim

309120-01

Boehringer Ingelheim Vetmedica, Inc.
2621 North Belt Highway • St. Joseph, MO 64506-2002
Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717
www.bi-vetmedica.com

BI S130931-RP-5 6/07