

Ingelvac® CircoFLEX®

Porcine Circovirus Vaccine Type 2, Killed Baculovirus Vector

Description

Ingelvac® CircoFLEX® is a single dose PCV2 vaccine for protection against circovirus in swine.

Features

For pigs 3 weeks of age and older

Benefits

- Vaccination at weaning is possible
- Protection before early exposure in nursery

Onset of immunity within 2 weeks of vaccination

Benefit

- Protection before early exposure in nursery

At least 17 weeks Duration of Immunity

Benefit

- Protection from PCVAD through the critical period in Grow-Finish

Single, 1 mL dose

Benefits

- Less labor costs than 2-dose vaccines
- Less stress on pigs
- Earlier onset of immunity than 2-dose vaccines

Highly syringeable

Benefit

- Less stress on labor force

No local or systemic side effects recorded

Benefits

- Less stress on pigs and no production set-back by vaccine
- Less chance of rejected pork at packing plant

21-day withdrawal period

Benefit

- Less risk of noncompliant pork entering food chain

Advantages

Ingelvac CircoFLEX is the only single-dose PCV2 vaccine that can be administered to pigs 3 weeks of age and older while providing at least 17 weeks DOI.

Ingelvac CircoFLEX contains Purified Circovirus Antigen (PCA™). PCA™ generates strong immunity and an excellent safety profile.

Dosage & Administration

Shake well before use. Using aseptic technique, administer a single 1 mL dose by intramuscular injection.

Precautions

Store in dark at 35-45° F (2-7° C). Do not freeze. Use entire contents when first opened.

Highlights

- For pigs as young as 3 weeks of age
- At least 17 weeks DOI
- PCV2 vaccine that is easy on pigs
- Contains PCA™

Uses


Recommended for the vaccination of healthy, susceptible pigs 3 weeks of age or older as an aid in the reduction of lymphoid depletion, inflammation and colonization of lymphoid tissue, viremia, and nasal shedding of virus associated with Porcine Circovirus Type 2 (PCV2).

Warning

Do not vaccinate within 21 days before slaughter. In case of anaphylactic reaction, epinephrine is symptomatic treatment.

Product Information

Name	Code	Size
Ingelvac®	130-931	50 dose
CircoFLEX™		50 mL
	130-941	100 dose
		100 mL
	130-951	250 dose
		250 mL



Exp. Date:
Serial No.:

Indications: For vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the prevention of lymphoid depletion, inflammation and colonization of lymphoid tissue associated with Porcine Circovirus Type 2 (PCV2). The onset of protection is 2 weeks post-vaccination. The duration of immunity is at least 17 weeks.

Directions and Dosage: See carton.

Precautions: Store at 35-45°F (2-7°C). Do not freeze. Shake well before using. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactic reaction, epinephrine is symptomatic treatment.

Preservative: Neomycin.



Porcine Circovirus Vaccine

Type 2, Killed Baculovirus Vector

Ingelvac CircoFLEX®

250 Doses/250 mL

Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, MO 64506
U.S. Veterinary License No. 124



Boehringer
Ingelheim

309120-01

Boehringer Ingelheim Vetmedica, Inc.
2621 North Belt Highway • St. Joseph, MO 64506-2002
Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717
www.bi-vetmedica.com

BI S130931-RP-5R 8/08