

Enterisol® SC-54 Enterisol® SC-54 FF

Salmonella choleraesuis Vaccine, Avirulent Live Culture

Description

Enterisol® SC-54 is recommended for use in healthy, susceptible, swine, one day of age or older, as an aid in the prevention of salmonellosis.

Features

Administration intranasally or via drinking water.

Safe for pigs one day of age or older.

Single dose administration.

Benefits

Fastest onset of immunity – 14 days.

Longest duration of immunity – at least 140 days.

Convenient administration causes less stress on pigs.

Saves labor costs with drinking water administration.

Reduces the prevalence of salmonellosis.

Advantages

Safety – The only salmonella vaccine that does not contain the virulence plasmid (disease causing gene).

The only vaccine proven to reduce salmonella in the live animal and at harvest.

Dosage & Administration

Lyophilized Form – Rehydrate the vaccine by adding the full contents of the accompanying sterile diluent to the vaccine. Shake well and use immediately.

Frozen Form – Thaw appropriate doses under slowly running cold tap water. Do not use warm water. Approximately 30 minutes per bottle.

Only the **lyophilized form** may be administered **intranasally** using a 2 mL dose for swine 1 day of age or older.

The **lyophilized form and the frozen form** may be administered **via the drinking water**. Administer via conventional (open trough or barrel-type tank system) or automatic watering systems equipped with proportioner. **See insert for complete directions.**

Precautions

Lyophilized form – Store at 35° - 45° F (2° - 7° C). Do not freeze.

Frozen form – Store with ice in conventional freezer for up to 10 days.

Uses

Mass vaccination to prevent salmonellosis.

Recommended for use in salmonella reduction programs.

Farms where pork quality is high priority.

Warning

Do not vaccinate within 21 days before slaughter. Use entire contents when first opened. Do not store reconstituted vaccine. In case of anaphylactoid reactions, epinephrine is symptomatic treatment.

Product Information

Name	Code	Size
Enterisol® SC-54	172-321	50 doses 100 mL
Enterisol® SC-54 FF	174-211	250 doses 100 mL
	174-221	500 doses 100 mL

Highlights

- Proven to reduce salmonella.
- Intranasal or water administration for lyophilized form.
- Water administration for frozen form.
- Avirulent live culture vaccine.
- Safe for use in pigs one day of age or older.



Serial No.:
Exp. Date:

Indications: For intranasal or water administration in healthy swine as an aid in the prevention of salmonellosis caused by *Salmonella choleraesuis* var. *kunzendorf*.

Directions: Reconstitute vaccine with accompanying diluent. Shake well. Place cannula on syringe and give 2 mL intranasally. See insert for water administration.

Precautions: Store at 35-45°F (2-7°C). Do not freeze. All materials used in administering this vaccine must be free of antibiotic or disinfectant residue to prevent inactivation. Avoid direct contact. If accidental human exposure occurs flush area with clean water and consult physician.

Salmonella Choleraesuis Vaccine

Avirulent Live Culture

Enterisol SC-54®

50 Doses/100 mL

Caution: Use entire contents when first opened. Burn this container and all unused contents. For Veterinary Use Only. Do not vaccinate within 21 days of slaughter.

Boehringer Ingelheim Vetmedica, Inc.

St. Joseph, MO 64506

(800) 325-9167

U.S. Veterinary License No. 124

U.S. Patent No. 5,436,001

472301N-00



**Boehringer
Ingelheim**

Serial No.:
Doses:
Exp. Date:

Salmonella Choleraesuis Vaccine, Avirulent Live Culture

Enterisol SC-54®FF

mL

Indications: For water administration in healthy swine 3 weeks of age or older as an aid in the prevention of salmonellosis caused by *Salmonella choleraesuis* var. *kunzendorf*.

Dosage: Administer 1 dose per pig in the drinking water. See insert for complete directions.

Precautions: Store frozen at ultra low temperatures below -50 degree C or with dry ice packs. Can be stored at -20 degrees C for no more than 14 days. Do not re-freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Burn containers and all unused contents. Anaphylactoid reactions may occur. Administer epinephrine. All materials used in administering this vaccine must be free of antimicrobial or disinfectant residue to prevent inactivation. Avoid contact. If accidental human exposure occurs, flush area with clean water & consult physician immediately. For veterinary use only.

Read entire insert before use.

Read the product label before use.

Boehringer Ingelheim Vetmedica, Inc.

St. Joseph, MO 64506

800-325-9167

U.S. Veterinary License No. 124

U.S. Patent No. 5,436,001

474201N-00



**Boehringer
Ingelheim**

Boehringer Ingelheim Vetmedica, Inc.

2621 North Belt Highway • St. Joseph, MO 64506-2002

Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717

www.bi-vetmedica.com