

Serial No.:
Exp. Date:

See carton for complete information.

Indications: For the vaccination of all types and ages of healthy horses against Eastern and Western Equine Encephalomyelitis, tetanus and as an aid in the prevention of Equine Influenza due to strains A₁ and A₂ viruses.

Administration and Dosage: Shake well. Inject 2 mL deep intramuscularly with a 1½ inch sterile needle preferably in the hind quarters. Revaccination required. See carton for full directions.

Caution: Store at 35-45°F (2-7°C). Protect from freezing. Use entire con-

**Encephalomyelitis-
Influenza Vaccine**

Eastern and Western,
Killed Virus

Tetanus Toxoid

Equi-Flu® EWT

10 Doses/20 mL

tents when first opened. Contains gentamicin and thimerosal as preservatives. Do not vaccinate within 21 days of slaughter. Anaphylactoid reactions may occur following use.

Antidote: Epinephrine.

Boehringer Ingelheim Vetmedica, Inc.

St. Joseph, MO 64506 U.S.A.

U.S. Veterinary License No. 124

Code 131-525; 31503-02



**Boehringer
Ingelheim**

Boehringer Ingelheim Vetmedica, Inc.

2621 North Belt Highway • St. Joseph, MO 64506-2002

Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717

www.bi-vetmedica.com