

Equi-Flu[®] EWT

Encephalomyelitis-Influenza Vaccine Eastern and Western, Killed Virus, Tetanus Toxoid

Description

Encephalomyelitis-Influenza Vaccine Eastern and Western, Killed Virus, Tetanus Toxoid. Equi-Flu[®] EWT contains gentamicin and thimerosal as preservatives.

Features

Provides broad spectrum active immunity against Eastern and Western Equine Encephalomyelitis.

Provides protection against relevant strains of equine influenza-A₁ and A₂ viruses.

Contains killed viruses and tetanus toxoid.

Benefits

Safe to use on all healthy horses.

Post-vaccination reactivity less than 1%.

Exceptionally non-irritating at injection site.

Advantages

Equi-Flu EWT is a killed virus vaccine that protects against Eastern and Western Encephalomyelitis, influenza, and tetanus.

Dosage & Administration

Using aseptic technique inject 2 mL deep IM with a 1½ inch sterile needle preferably in the hind quarters. For primary vaccination administer a second 2 mL dose in 4 to 6 weeks using a different injection site. Revaccinate annually for protection against tetanus, using a single 2 mL dose. Revaccinate annually and prior to anticipated exposure for maximum benefit against Equine Influenza strains A₁ and A₂ viruses and Eastern and Western Equine Encephalomyelitis.

Precautions

Store out of direct sunlight at 35° - 45° F (2° - 7° C). Shake well. Protect from freezing. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur following use. Antidote: Epinephrine.

Uses

Equi-Flu EWT is recommended for vaccination of all types and ages of healthy horses against Eastern and Western Equine Encephalomyelitis, tetanus and as an aid in prevention of Equine Influenza due to strains A₁ and A₂ viruses.

Warning

This product has been tested under laboratory conditions and shown to meet all Federal standards for safety and efficacy in normal healthy animals. This level of performance may be affected by conditions such as stress, weather, nutrition, disease, parasitism, other treatments, individual idiosyncrasies or impaired immunological competency. These factors should be considered by the user when evaluating product performance or freedom from reactions.

Product Information

Name	Code	Size
Equi-Flu [®] EWT	131-515	2 mL 1 dose
	131-525	20 mL 10 dose

Highlights

- Non-irritating injection.
- Sold only to licensed veterinarians.



Serial No.:
Exp. Date:

See carton for complete information.

Indications: For the vaccination of all types and ages of healthy horses against Eastern and Western Equine Encephalomyelitis, tetanus and as an aid in the prevention of Equine Influenza due to strains A₁ and A₂ viruses.

Administration and Dosage: Shake well. Inject 2 mL deep intramuscularly with a 1½ inch sterile needle preferably in the hind quarters. Revaccination required. See carton for full directions.

Caution: Store at 35-45°F (2-7°C). Protect from freezing. Use entire con-

**Encephalomyelitis-
Influenza Vaccine**

Eastern and Western,
Killed Virus

Tetanus Toxoid

Equi-Flu® EWT

10 Doses/20 mL

tents when first opened. Contains gentamicin and thimerosal as preservatives. Do not vaccinate within 21 days of slaughter. Anaphylactoid reactions may occur following use.

Antidote: Epinephrine.

Boehringer Ingelheim Vetmedica, Inc.

St. Joseph, MO 64506 U.S.A.

U.S. Veterinary License No. 124

Code 131-525; 31503-02



**Boehringer
Ingelheim**

Boehringer Ingelheim Vetmedica, Inc.

2621 North Belt Highway • St. Joseph, MO 64506-2002

Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717

www.bi-vetmedica.com