

Serial No.:  
Exp. Date:

**Indications:** For the vaccination of all types and ages of healthy horses against Eastern, Western and Venezuelan Equine Encephalomyelitis, tetanus, and as an aid in the prevention of Equine Influenza due to strains A<sub>1</sub> and A<sub>2</sub> viruses.

**Administration and Dosage:** Shake well. Inject 2 mL deep intramuscularly with a 1½ inch sterile needle preferably in the hind quarters. Revaccination required. See carton for full directions.

**Caution:** Store at 35-45°F (2-7°C). Protect from freezing. Use entire contents when first opened. Contains gen-

## Encephalomyelitis- Influenza Vaccine

Eastern, Western, and  
Venezuelan, Killed Virus,

## Tetanus Toxoid

Equi-Flu® VEWT

10 Doses/20 mL

tamicin and thimerosal as preservatives. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur following use.

**Antidote:** Epinephrine.

Boehringer Ingelheim Vetmedica, Inc.  
St. Joseph, MO 64506 U.S.A.  
U.S. Veterinary License No. 124

Code 131-625; 31603-02



Boehringer  
Ingelheim

**Boehringer Ingelheim Vetmedica, Inc.**

**2621 North Belt Highway • St. Joseph, MO 64506-2002**

**Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717**

**www.bi-vetmedica.com**