

# Equi-Flu<sup>®</sup> VEWT

## Encephalomyelitis-Influenza Vaccine Eastern, Western, and Venezuelan, Killed Virus Tetanus Toxoid

### Description

Encephalomyelitis-Influenza Vaccine Eastern, Western, and Venezuelan, Killed Virus. Equi-Flu<sup>®</sup> VEWT contains gentamicin and thimerosal as preservatives.

### Features

Provides broad spectrum active immunity against Eastern, Western, and Venezuelan Equine Encephalomyelitis.

Provides protection against relevant strains of Equine Influenza-A<sub>1</sub> and A<sub>2</sub> viruses.

Contains killed viruses and tetanus toxoid.

### Benefits

Safe to use on all healthy horses.

Exceptionally non-irritating at injection site.

Post-vaccination reactivity less than 1%.

### Advantages

Equi-Flu VEWT is a killed virus vaccine that protects against Eastern, Western, and Venezuelan Equine Encephalomyelitis, influenza, and tetanus.

### Dosage & Administration

Using aseptic technique inject 2 mL deep IM with a 1½ inch sterile needle preferably in the hind quarters. For primary vaccination administer a second 2 mL dose in 4 to 6 weeks using a different injection site. Revaccinate annually for protection against tetanus, using a single 2 mL dose. Revaccinate annually and prior to anticipated exposure for maximum benefit against Equine Influenza strains A<sub>1</sub> and A<sub>2</sub> viruses and Eastern, Western, and Venezuelan Equine Encephalomyelitis.

### Precautions

Store out of direct sunlight at 35° - 45° F (2° - 7° C). Shake well. Protect from freezing. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur following use. Antidote: Epinephrine.

### Uses

Equi-Flu VEWT is recommended for vaccination of all types and ages of healthy horses against Eastern, Western, and Venezuelan Equine Encephalomyelitis, tetanus and as an aid in prevention of Equine Influenza strains A<sub>1</sub> and A<sub>2</sub>.

### Warning

This product has been tested under laboratory conditions and shown to meet all Federal standards for safety and efficacy in normal healthy animals. This level of performance may be affected by conditions such as stress, weather, nutrition, disease, parasitism, other treatments, individual idiosyncrasies or impaired immunological competency. These factors should be considered by the user when evaluating product performance or freedom from reactions.

### Product Information

Name	Code	Size
Equi-Flu <sup>®</sup> VEWT	131-615	2 mL 1 dose
	131-625	20 mL 10 dose

### Highlights

- Non-irritating.
- Sold only to licensed veterinarians.



Serial No.:  
Exp. Date:

**Indications:** For the vaccination of all types and ages of healthy horses against Eastern, Western and Venezuelan Equine Encephalomyelitis, tetanus, and as an aid in the prevention of Equine Influenza due to strains A<sub>1</sub> and A<sub>2</sub> viruses.

**Administration and Dosage:** Shake well. Inject 2 mL deep intramuscularly with a 1½ inch sterile needle preferably in the hind quarters. Revaccination required. See carton for full directions.

**Caution:** Store at 35-45°F (2-7°C). Protect from freezing. Use entire contents when first opened. Contains gen-

## Encephalomyelitis- Influenza Vaccine

Eastern, Western, and  
Venezuelan, Killed Virus,

## Tetanus Toxoid

Equi-Flu® VEWT

10 Doses/20 mL

tamicin and thimerosal as preservatives. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur following use.

**Antidote:** Epinephrine.

Boehringer Ingelheim Vetmedica, Inc.  
St. Joseph, MO 64506 U.S.A.

U.S. Veterinary License No. 124

Code 131-625; 31603-02



Boehringer  
Ingelheim

**Boehringer Ingelheim Vetmedica, Inc.**

**2621 North Belt Highway • St. Joseph, MO 64506-2002**

**Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717**

**www.bi-vetmedica.com**