

Tetguard™

Tetanus Toxoid

Description

Tetanus Toxoid. Tetguard™ is alum precipitated, purified for vaccination against tetanus. Contains thimerosal as a preservative.

Features

Approved for use in horses, cattle, sheep, and swine.

Provides long-term, active immunity against tetanus.

Sub-Q and IM administration.

Benefits

Special treatment eliminates toxicity and retains antigenicity.

Advantages

Tetguard is manufactured by a special treatment method that modifies the virulent tetanus bacilli to eliminate toxicity while retaining antigenicity.

Dosage & Administration

Using aseptic technique administer Sub-Q or IM.

Horses, cattle, swine: Administer 1 mL.

Sheep: Administer 0.5 mL. Repeat in 30 days. Revaccinate annually or prior to anticipated exposure.

Precautions

Store out of direct sunlight at 35° - 45° F (2° - 7° C). Shake bottle well. Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur following use.

Antidote: Epinephrine

Uses

Tetguard is recommended for vaccination of healthy horses, cattle, swine, and sheep to confer long-term, active immunity against tetanus.

Warning

This product has been tested under laboratory conditions and shown to meet all Federal standards for safety and efficacy in normal healthy animals. This level of performance may be affected by conditions such as stress, weather, nutrition, disease, parasitism, other treatments, individual idiosyncrasies or impaired immunological competency. These factors should be considered by the user when evaluating product performance or freedom from reactions.

Highlights

- Special treatment eliminates toxicity while retaining antigenicity.
- Sold only to licensed veterinarians.
- Approved for use in horses, cattle, sheep, swine.

Product Information

Name	Code	Size
Tetguard™	131-815	1 mL
	131-825	20 mL




Indications: Alum precipitated, purified toxoid for vaccination against tetanus.

Administration and Dosage: Shake well before use. Administer subcutaneously or intramuscularly, using aseptic technique.
Dosage: Horses, cattle and swine – 1 mL. Sheep – 0.5 mL. Repeat in 30 days. Revaccinate annually and prior to anticipated exposure.

Serial No.:

Exp. Date:



Boehringer Ingelheim
Vetmedica, Inc.
St. Joseph, MO 64506 U.S.A.
U.S. Veterinary License No. 124
Code 131-815; 31801-03



Caution: Store at 35-45°F (2-7°C). Protect from freezing. Use entire contents when first opened. Contains thimerosal as a preservative. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur following use.

Antidote: Epinephrine.

Precaution: This product has been tested under laboratory conditions and shown to meet all Federal standards for safety and efficacy in normal healthy animals. This level of performance may be affected by conditions such as stress, weather, nutrition, disease, parasitism, other treatments, individual idiosyncrasies or impaired immunological competency. These factors should be considered by the user when evaluating product performance or freedom from reactions.



Boehringer Ingelheim Vetmedica, Inc.
2621 North Belt Highway • St. Joseph, MO 64506-2002
Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717
www.bi-vetmedica.com