




Bio-Mycin[®] 200 Head-to-Head Study Results.

 <p>Bio-Mycin[®] 200 (Oxytetracycline)</p>	<p>Injection site reaction has resolved</p>
 <p>Liquamycin[®] LA-200[®] or generics</p>	<p>Injection site reaction is mild to moderate</p>
 <p>Tetradure[™] 300 (Noromycin[®] 300 LA)</p>	<p>Injection site is moderate to severe and unresolved or still active</p>

In a head-to-head study, Bio-Mycin[®] 200 (Sub-Q or IM) showed reduced tissue damage in cattle 48 days after injection when compared to Nuflor[®], Micotil[®], Liquamycin[®] LA-200[®] (Sub-Q or IM) and Tetradure[™] 300.¹

Bio-Mycin[®] 200 is approved for lactating and non-lactating dairy cows and all beef cattle.

WARNING: Discontinue treatment at least 28 days prior to slaughter of cattle and swine. Milk taken from animals during treatment and for 96 hours after the last treatment must not be used for food. Rapid intravenous administration may result in animal collapse. Oxytetracycline should be administered intravenously slowly over a period of at least 5 minutes.

¹Dowling P. Evaluation of SQ and IM injection sites of antimicrobials in calves. University of Saskatchewan.

Bio-Mycin is a registered trademark of Boehringer Ingelheim Vetmedica, Inc. Liquamycin and LA-200 are registered trademarks of Zoetis Inc., its affiliates and/or its licensors. Micotil is a registered trademark for Elanco's brand of tilmicosin injection. Nuflor is a registered trademark of Intervet/Schering-Plough Animal Health. Noromycin 300 LA is a registered trademark of Norbrook Laboratories Limited. Tetradure is a trademark of Merial.
© 2018 Boehringer Ingelheim Vetmedica, Inc. BOV-1373-ANTB0618