

Exp. Date:

Serial No.:



Indications: Recommended for the vaccination of healthy, susceptible sheep and cattle, including bred and/or lactating beef and dairy cattle, as an aid in the reduction of diseases caused by *Clostridium chauvoei*, *Cl. septicum*, *Cl. novyi*, *Cl. sordellii*, and *Cl. perfringens* Types C and D. Although *Cl. perfringens* Type B is not a significant problem in the U.S.A., immunity may be provided against the beta and epsilon toxins elaborated by *Cl. perfringens* Type B. This immunity is derived from the combination of Type C (beta) and Type D (epsilon) fractions.

Dosage: Using aseptic technique, inject 2 mL subcutaneously.

**Clostridium
Chauvoei-Septicum-
Novyi-Sordellii-
Perfringens Types C&D
Bacterin-Toxoid**

Caliber™ 7

50 Doses/100 mL

Repeat in 21 to 28 days and once annually.

Precautions: Store at 35-45°F (2-7°C). Avoid freezing. Shake well before using. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur.

Antidote: Epinephrine.

**Boehringer Ingelheim
Vetmedica, Inc.**

St. Joseph, MO 64506 U.S.A.

U.S. Veterinary License No. 124

Code 133-331; 33302-01



**Boehringer
Ingelheim**

Boehringer Ingelheim Vetmedica, Inc.

2621 North Belt Highway • St. Joseph, MO 64506-2002

Information 800-325-9167 • Orders 800-325-9167 • Fax 816-236-2717

www.bi-vetmedica.com