

Serial No.:  
Exp. Date:

## Lawsonia Intracellularis Vaccine

Avirulent Live Culture

### Enterisol® Ileitis FF

For use in swine only

100 Doses/100 mL

**Indications:** Recommended for the vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the prevention and control of porcine proliferative enteropathy (ileitis) caused by *Lawsonia intracellularis*. In clinical studies the vaccine significantly reduced gross and microscopic intestinal lesions of ileitis. Vaccinated pigs also had significantly reduced colonization by *Lawsonia intracellularis* following challenge with the virulent microorganism. Protective immunity following vaccination has been demonstrated to be at least 7 weeks. Enterisol® Ileitis FF may be simultaneously administered orally to pigs along with Enterisol® SC-54 FF and/or Ingelvac® ERY-ALC.

**Dosage:** Administer 1 dose per pig in the drinking water, to pigs 3 weeks of age or older. See insert for complete directions.

**Precautions:** Store frozen at -70°C. Do not thaw until immediately prior to use. Shake well before use. Do not vaccinate within 21 days before slaughter. Burn container and all unused contents. Do not use in pregnant swine or breeding boars. Do not use concurrently with antibiotics, including feed antibiotics. A *minimum* antibiotic-free period of 3 days pre- and post-vaccination is recommended. All materials used in administering this vaccine must be free of antimicrobial or disinfectant residue to prevent inactivation. Anaphylactic reactions may occur.

**Antidote:** Epinephrine.

Boehringer Ingelheim Vetmedica, Inc. / St. Joseph, MO 64506  
U.S. Veterinary License No. 124 / U.S. Patent Nos. 5,714,375 and 5,885,823  
Code 129-641



Boehringer  
Ingelheim

29611-01

100 Doses

Serial No.:  
Exp. Date:

## Lawsonia Intracellularis Vaccine

Avirulent Live Culture

### Enterisol® Ileitis FF

For use in swine only

250 Doses/100 mL

**Indications:** Recommended for the vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the prevention and control of porcine proliferative enteropathy (ileitis) caused by *Lawsonia intracellularis*. In clinical studies the vaccine significantly reduced gross and microscopic intestinal lesions of ileitis. Vaccinated pigs also had significantly reduced colonization by *Lawsonia intracellularis* following challenge with the virulent microorganism. Protective immunity following vaccination has been demonstrated to be at least 7 weeks. Enterisol® Ileitis FF may be simultaneously administered orally to pigs along with Enterisol® SC-54 FF and/or Ingelvac® ERY-ALC.

**Dosage:** Administer 1 dose per pig in the drinking water, to pigs 3 weeks of age or older. See insert for complete directions.

**Precautions:** Store frozen at -70°C. Do not thaw until immediately prior to use. Shake well before use. Do not vaccinate within 21 days before slaughter. Burn container and all unused contents. Do not use in pregnant swine or breeding boars. Do not use concurrently with antibiotics, including feed antibiotics. A *minimum* antibiotic-free period of 3 days pre- and post-vaccination is recommended. All materials used in administering this vaccine must be free of antimicrobial or disinfectant residue to prevent inactivation. Anaphylactic reactions may occur.

**Antidote:** Epinephrine.

Boehringer Ingelheim Vetmedica, Inc. / St. Joseph, MO 64506  
U.S. Veterinary License No. 124 / U.S. Patent Nos. 5,714,375 and 5,885,823  
Code 129-671



Boehringer  
Ingelheim

29612-01

250 Doses

Serial No.:  
Exp. Date:

## Lawsonia Intracellularis Vaccine

Avirulent Live Culture

### Enterisol® Ileitis FF

For use in swine only

500 Doses/100 mL

**Indications:** Recommended for the vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the prevention and control of porcine proliferative enteropathy (ileitis) caused by *Lawsonia intracellularis*. In clinical studies the vaccine significantly reduced gross and microscopic intestinal lesions of ileitis. Vaccinated pigs also had significantly reduced colonization by *Lawsonia intracellularis* following challenge with the virulent microorganism. Protective immunity following vaccination has been demonstrated to be at least 7 weeks. Enterisol® Ileitis FF may be simultaneously administered orally to pigs along with Enterisol® SC-54 FF and/or Ingelvac® ERY-ALC.

**Dosage:** Administer 1 dose per pig in the drinking water, to pigs 3 weeks of age or older. See insert for complete directions.

**Precautions:** Store frozen at -70°C. Do not thaw until immediately prior to use. Shake well before use. Do not vaccinate within 21 days before slaughter. Burn container and all unused contents. Do not use in pregnant swine or breeding boars. Do not use concurrently with antibiotics, including feed antibiotics. A *minimum* antibiotic-free period of 3 days pre- and post-vaccination is recommended. All materials used in administering this vaccine must be free of antimicrobial or disinfectant residue to prevent inactivation. Anaphylactic reactions may occur.

**Antidote:** Epinephrine.

Boehringer Ingelheim Vetmedica, Inc. / St. Joseph, MO 64506  
U.S. Veterinary License No. 124 / U.S. Patent Nos. 5,714,375 and 5,885,823  
Code 129-681



Boehringer  
Ingelheim

29613-01

500 Doses