

Exp. Date:

Serial No.:

Indications: Recommended for the vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the reduction of enzootic pneumonia of swine caused by *Mycoplasma hyopneumoniae*.

Composition: Prepared from chemically inactivated cultures of *M. hyopneumoniae*, combined with an adjuvant to stimulate the immune response.

Directions and Dosage: Shake well before use. Using aseptic technique, inject a single 1 mL dose intramuscularly. The duration of immunity is at least 26 weeks. Semi-annual revaccination is recommended for breeding swine.



Mycoplasma Hyopneumoniae Bacterin

Ingelvac
MycoFLEX[®]

50 Doses/50 mL

Precautions: Store at 35–45°F (2–7°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur. Antidote: Epinephrine. **Caution: Do not use as diluent for live vaccines.**

Boehringer Ingelheim
Vetmedica, Inc.
St. Joseph, MO 64506
U.S. Veterinary License No. 124



Boehringer
Ingelheim

27301-01

Exp. Date:

Serial No.:

Indications: Recommended for the vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the reduction of enzootic pneumonia of swine caused by *Mycoplasma hyopneumoniae*.

Composition: Prepared from chemically inactivated cultures of *M. hyopneumoniae*, combined with an adjuvant to stimulate the immune response.

Directions and Dosage: Shake well before use. Using aseptic technique, inject a single 1 mL dose intramuscularly. The duration of immunity is at least 26 weeks. Semi-annual revaccination is recommended for breeding swine.

Precautions: Store at 35–45°F (2–7°C). Do not freeze. Use entire contents when first



Mycoplasma Hyopneumoniae Bacterin

Ingelvac
MycoFLEX[®]

100 Doses/100 mL

opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur. Antidote: Epinephrine.
Caution: Do not use as diluent for live vaccines.

Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, MO 64506
U.S. Veterinary License No. 124



Boehringer
Ingelheim

27302-01

Serial No.:
Exp. Date:

Indications: Recommended for the vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the reduction of enzootic pneumonia of swine caused by *Mycoplasma hyopneumoniae*.

Composition: Prepared from chemically inactivated cultures of *M. hyopneumoniae*, combined with an adjuvant to stimulate the immune response.

Directions and Dosage: Shake well before use. Using aseptic technique, inject a single 1 mL dose intramuscularly. The duration of immunity is at least 26 weeks. Semi-annual revaccination is recommended for breeding swine.

Precautions: Store at 35–45°F (2–7°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur. Antidote: Epinephrine. **Caution:** Do not use as diluent for live vaccines.



Mycoplasma Hyopneumoniae Bacterin

Ingelvac MycoFLEX[®]

250 Doses/250 mL

Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, MO 64506
U.S. Veterinary License No. 124



Boehringer
Ingelheim

27303-01