

time of stress or exposure. Calves vaccinated under six months of age should be revaccinated at six months of age. May be administered to pregnant animals at any stage of gestation. Protect animals from exposure for at least 14 days after the last dose of vaccine.

Precautions: Store in the dark at 35 - 45°F (2 - 7°C). Avoid freezing. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine.

**Boehringer Ingelheim
Vetmedica, Inc.**
St. Joseph, MO 64506
U.S. Vet. License No. 124
Code 161-211



**Boehringer
Ingelheim**

**Bovine Rhinotracheitis-Virus
Diarrhea-Parainfluenza 3-
Respiratory Syncytial Virus
Vaccine, Killed Virus, *Leptospira
Canicola-Grippotyphosa-
Hardjo-Icterohaemorrhagiae-
Pomona Bacterin***

TRIANGLE[®] 10

10 Doses/50 mL

Indications: For vaccination of healthy cattle as an aid in the prevention of disease caused by infectious bovine rhinotracheitis (IBR), bovine virus diarrhea (BVD types 1 & 2), parainfluenza 3 (PI3), and bovine respiratory syncytial (BRSV) viruses, *Leptospira pomona*, *L. hardjo*, *L. grippotyphosa*, *L. canicola*, and *L. icterohaemorrhagiae*.

Composition: Contains IBR, BVD Types 1 and 2, PI3, and BRSV killed viruses, *L. pomona*, *L. hardjo*, *L. grippotyphosa*, *L. canicola*, and *L. icterohaemorrhagiae*, with neomycin, polymyxin B, and thimerosal as preservatives.

Dosage and Administration: Shake well. Inject one 5 mL dose intramuscularly or subcutaneously using aseptic technique. Repeat in 14 to 28 days. A 5 mL booster dose is recommended annually or prior to



Serial No.:

Exp. Date:

Serial No.:
Exp. Date:



Indications: For vaccination of healthy cattle as an aid in the prevention of disease caused by infectious bovine rhinotracheitis (IBR), bovine virus diarrhea (BVD types 1 & 2), parainfluenza 3 (PI3), and bovine respiratory syncytial (BRSV) viruses, *Leptospira pomona*, *L. hardjo*, *L. grippotyphosa*, *L. canicola*, and *L. icterohaemorrhagiae*.

Composition: Contains IBR, BVD Types 1 and 2, PI3, and BRSV killed viruses, *L. pomona*, *L. hardjo*, *L. grippotyphosa*, *L. canicola*, and *L. icterohaemorrhagiae*, with neomycin, polymyxin B, and thimerosal as preservatives.

Dosage and Administration: Shake well. Inject one 5 mL dose intramuscularly or subcutaneously using aseptic technique. Repeat in 14 to 28 days. A 5 mL booster dose is

**Bovine Rhinotracheitis-Virus
Diarrhea-Parainfluenza 3-
Respiratory Syncytial Virus
Vaccine, Killed Virus, Leptospira
Canicola-Grippytyphosa-
Hardjo-Icterohaemorrhagiae-
Pomona Bacterin**

TRIANGLE[®] 10

50 Doses/250 mL

recommended annually or prior to time of stress or exposure. Calves vaccinated under six months of age should be revaccinated at six months of age. May be administered to pregnant animals at any stage of gestation. Protect animals from exposure for at least 14 days after the last dose of vaccine.

Precautions: Store in the dark at 35 - 45°F (2 - 7°C). Avoid freezing. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine.

Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, MO 64506
U.S. Vet. License No. 124
Code 161-231



**Boehringer
Ingelheim**

61202-01