

Boehringer
Ingelheim

Revision 1
Operator hal
Date 24Feb11
Art # 60802-01
Item # 460801
Product Triangle 5

Type/Size 50 Dose Label
Template # 289
Dimensions 2" x 5"
File Path V1/BI Labeling
PDF..... 21Feb11 hal
Rev Level 01

BLACK

PMS 362

PMS 205

VARNISH

SEPARATIONS
CHECKED by

hal 24Feb11

Click here with mover tool to bring
varnish plate to front

Click here with mover tool to bring keyline to front

Serial No.:
Exp. Date:



Indications: For vaccination of healthy cattle as an aid in the prevention of disease caused by infectious bovine rhinotracheitis (IBR), bovine virus diarrhea (BVD types 1 & 2), parainfluenza 3 (PI3), and bovine respiratory syncytial (BRSV) viruses.

Composition: Contains IBR, BVD Types 1 and 2, PI3, and BRSV killed viruses, with neomycin, polymyxin B, and thimerosal as preservatives.

Dosage and Administration: Shake well before using. Inject one 2 mL dose intramuscularly or subcutaneously, using aseptic technique. Repeat in 14 to 28 days. A 2 mL booster dose is recommended annually or prior to time of stress or exposure. Calves vaccinated under six months of age should be revaccinated at six months of age. May be administered to pregnant animals at

Bovine Rhinotracheitis- Virus Diarrhea- Parainfluenza 3- Respiratory Syncytial Virus Vaccine, Killed Virus

TRIANGLE[®] 5

50 Doses/100 mL

any stage of gestation. Protect animals from exposure for at least 14 days after the last dose of vaccine.

Precautions: Store in the dark at 35 - 45°F (2 - 7°C). Avoid freezing. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Transient swelling may occur at the injection site. In case of anaphylactoid reaction, administer epinephrine.

Boehringer Ingelheim
Vetmedica, Inc.
St. Joseph, MO 64506
U.S. Veterinary License No. 124
Code: 160-831



Boehringer
Ingelheim

60802-01