

**Boehringer
Ingelheim**

Revision ... 2

Operator ... lsh

Date 13Feb14

Art# 161601-01

Item # 461600

SAP # 107314-002

Product ... TrichGuard V5L

Type/Size Label / 10 Doses, 50 mL

Template # 290

Dimensions 1-3/4" x 4"

Corner Radius 0.94"

PDF 13Feb14 lsh

Rev Level 01

MASTER

BLACK

PMS 151

PMS 873

VARNISH

[Click here with mover tool to bring varnish plate to front](#)

Serial No.:
Exp. Date:

0
1231316161
3

Indications: For vaccination of healthy cattle as an aid in the prevention of disease caused by *Campylobacter fetus*, *Leptospira canicola*, *L. grippityphosa*, *L. hardjo*, *L. icterohaemorrhagiae* and *L. pomona*, and as an aid in the reduction of shedding of *Tritrichomonas foetus*.

Composition: Killed, concentrated cultures of *Tritrichomonas foetus*, *Campylobacter fetus* var. *veneralis*, *Leptospira canicola*, *L. grippityphosa*, *L. hardjo*, *L. icterohaemorrhagiae*, and *L. pomona* organisms suspended in an oil adjuvant. Contains neomycin, polymyxin B, and thimerosal as preservatives.

Dosage and administration: Shake well before using. Inject one 5 mL dose subcutaneously under aseptic conditions, in the

**Trichomonas Foetus
Vaccine, Killed Protozoa
Campylobacter Fetus-
Leptospira Canicola-
Grippityphosa-Hardjo-
Icterohaemorrhagiae-
Pomona Bacterin**

TrichGuard® V5L
10 Doses/50 mL

neck or behind the shoulder. A second dose should be administered 2 to 4 weeks later. The last injection should precede the breeding season by 4 weeks. Revaccinate annually.

Precautions: Store in the dark at 35 - 45°F (2 - 7°C). Avoid freezing. Use entire contents when first opened. Do not vaccinate within 60 days before slaughter. Transient swelling may occur at injection site. In case of anaphylactoid reaction, administer epinephrine.

Boehringer Ingelheim
Vetmedica, Inc.
St. Joseph, MO 64506
US Vet. Lic. No. 124

 **Boehringer
Ingelheim**

161601-01