

Exp.:

Ser.:

See carton for complete information.

Indications: For vaccination of healthy horses 4 months of age or older as an aid in the prevention of disease caused by Eastern, Western, and Venezuelan Encephalomyelitis, and tetanus.

Precautions: Store at 35-45°F (2-7°C). Protect from freezing. Do not vaccinate within 21 days before slaughter. Use entire contents when first opened.

Encephalomyelitis Vaccine

Eastern & Western & Venezuelan, Killed Virus

Tetanus Toxoid

Vetera® VEWT

10 Doses/10 mL

Anaphylactoid reactions may occur following use. Antidote: Epinephrine.

Preservatives: Gentamicin and Amphotericin B.

For Veterinary Use Only

Boehringer Ingelheim Vetmedica, Inc.

St. Joseph, MO 64506

U.S. Veterinary License No. 124

Code 135-725



**Boehringer
Ingelheim**

135703-00

Vetera is a registered trademark of Boehringer Ingelheim Vetmedica, GmbH, Ingelheim, Germany

Carbimmune, North American Equine E159 and Ultrafil are trademarks of Boehringer Ingelheim Vetmedica, Inc.

Boehringer Ingelheim Vetmedica, Inc.
2621 North Belt Highway • St. Joseph, MO 64506-2002
Information 800-325-9167 • Orders 800-325-9167 • Fax 816-236-2717
www.bi-vetmedica.com