

Exp. Date:

Serial No.:



TAKE TIME

OBSERVE LABEL
DIRECTIONS

Indications: Recommended for the vaccination of healthy, susceptible cattle and sheep against diseases caused by *Clostridium chauvoei*, *Cl. septicum*, *Cl. haemolyticum*, *Cl. novyi*, *Cl. sordellii* and *Cl. perfringens* Types C and D. Although *Clostridium perfringens* Type B is not a significant problem in the U.S.A., immunity may be provided against the beta and epsilon toxins elaborated by *Clostridium perfringens* Type B. This immunity is derived from the combination of Type C (beta) and Type D (epsilon) fractions.

Dosage:

Cattle: Using aseptic technique, inject 5 mL subcutaneously. Repeat in 21 to 28 days and once annually.

Sheep: Using aseptic technique, inject 2.5 mL subcutaneously. Repeat in 21 to 28 days and once annually.

Clostridium Chauvoei- Septicum-Haemolyticum- Novyi-Sordellii-Perfringens Types C & D Bacterin-Toxoid

Bar-Vac® 8

(7-Way + Redwater)

250 mL/50 Cattle Doses
or 100 Sheep Doses

Precautions: Store out of direct sunlight at a temperature between 35-45°F (2-7°C). Avoid freezing. Shake well before using. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur.

Antidote: Epinephrine.

Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, Missouri 64506 U.S.A.
U.S. Veterinary License No. 124

Code 124-231; 24202-02



Boehringer
Ingelheim