

Serial No.:
Exp. Date:

Indications: Recommended for the vaccination of healthy, susceptible sheep and cattle against enterotoxemia caused by the toxins of *Clostridium perfringens* Types C and D. Although *Cl. perfringens* Type B is not a significant problem in the U.S.A., immunity may be provided against the beta and epsilon toxins elaborated by *Cl. perfringens* Type B. This immunity is derived from the combination of Type C (beta) and Type D (epsilon) fractions.

Composition: Prepared from cultures of *Clostridium perfringens* Types C and D. Alum Precipitated.

Dosage:

Sheep: Using aseptic technique, inject 2 mL subcutaneously. Repeat in 21 to 28 days and annually.

Cattle: Using aseptic technique,

Clostridium Perfringens Types C & D Toxoid

Bar-Vac® CD

250 mL/50 Cattle Doses
or 125 Sheep Doses

inject 5 mL subcutaneously. Repeat in 21 to 28 days and annually.

Precautions: Store out of direct sunlight at a temperature between 35-45°F (2-7°C). Avoid freezing. Shake well before using. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur.

Antidote: Administer epinephrine.

**Boehringer Ingelheim
Vetmedica, Inc.**
St. Joseph, Missouri 64506 U.S.A.
U.S. Veterinary License No. 124

Code 110-921; 10902-03



**Boehringer
Ingelheim**

Boehringer Ingelheim Vetmedica, Inc.
2621 North Belt Highway • St. Joseph, MO 64506-2002
Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2731
www.bi-vetmedica.com