

FLEXcombo®

Porcine Circovirus Vaccine Type 2, Killed Baculovirus Vector Mycoplasma Hypopneumoniae Bacterin

Description

The FLEXcombo® package contains one bottle of Ingelvac CircoFLEX® and one bottle of Ingelvac MycoFLEX®. The Ingelvac MycoFLEX® is supplied in a headspace bottle (HSB) with extra capacity for the purpose of convenient mixing with Ingelvac CircoFLEX®.

Advantages

- Ability to customize control of multiple diseases with a single injection
- Single, 2 mL dose
- Reduced labor costs compared to 2-dose vaccines
- Single dose administration minimizes stress on pigs
- Contains ImpranFLEX® adjuvant effective, aqueous-based (non-oil) polymer adjuvant
- PQA Plus® friendly
- Low viscosity, highly syringeable
- Easy administration
- Member of the FLEX Family™
- New headspace bottle (HSB) technology for convenient mixing

Benefits

Boehringer Ingelheim Vetmedica, Inc. research data demonstrate compatibility between Ingelvac CircoFLEX® and Ingelvac MycoFLEX®, offering convenient control of mycoplasma and circovirus in a single injection.

The ImpranFLEX® adjuvant utilized in FLEXcombo® allows rapid onset of immunity.

Low risk of adverse reactions.

Dosage & Administration

Shake each bottle before use. Aseptically mix the full contents of both bottles as per the mixing instruction diagram; see back side of this carton. Shake well. Inject a single 2 mL dose intramuscularly. Semi-annual revaccination with Ingelvac MycoFLEX is recommended for breeding swine.

Precautions

Store at 35-45° F (2-7° C).
Do not freeze.

Uses

For vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the prevention of lymphoid depletion, inflammation and colonization of lymphoid tissue associated with Porcine Circovirus Type 2 (PCV2), and as an aid in the prevention of disease caused by *Mycoplasma hyopneumoniae*. The onset of protection against Porcine Circovirus Type 2 is 2 weeks post-vaccination, and the duration of immunity is at least 4 months. The duration of immunity against *M. hyopneumoniae* is at least 26 weeks.

Warning

Do not vaccinate within 21 days before slaughter. In case of anaphylactic reaction, epinephrine is symptomatic treatment.

Product Information

Name	Code	Size
FLEXcombo®	100506	250 dose
	100505	50 dose

Highlights

- FLEXcombo® with headspace bottle technology for convenient mixing
- Highly syringeable
- Ability to customize vaccination scheduling
- Single 2 mL dose



Indications: For vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the prevention of lymphoid depletion, inflammation and colonization of lymphoid tissue associated with Porcine Circovirus Type 2 (PCV2), and as an aid in the prevention of disease caused by *Mycoplasma hyopneumoniae*. The onset of protection against Porcine Circovirus Type 2 is 2 weeks post-vaccination, and the duration of immunity is at least 4 months. The duration of immunity against *M. hyopneumoniae* is at least 26 weeks.

Directions and dosage: Shake each bottle before use. Aseptically mix the full contents of both bottles as per the mixing instruction diagram; see back side of this carton. Shake well. Inject a single 2 mL dose intramuscularly. Semi-annual revaccination with Ingelvac MycoFLEX is recommended for breeding swine.


Porcine Circovirus Vaccine
 Type 2, Killed Baculovirus Vector
Mycoplasma
Hypneumoniae Bacterin
FLEXcombo®
250 Doses HSB
 Mix to 500 mL

Precautions: Store at 35-45°F (2-7°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactic reaction, epinephrine is symptomatic treatment.

Preservative: Neomycin.
 This package contains one 250 dose (250 mL) vial of Ingelvac CircoFLEX® and one 250 dose (250 mL) vial of Ingelvac MycoFLEX®. The Ingelvac MycoFLEX is supplied in a headspace bottle (HSB) with extra capacity for the purpose of aseptic mixing.



Boehringer Ingelheim
 Vetmedica, Inc.
 St. Joseph, MO 64506
 US Vet. Lic. No. 124



Boehringer Ingelheim Vetmedica, Inc.
 3902 Genefield Road • St. Joseph, MO 64506
 Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717
www.bi-vetmedica.com