

Serial No.:
Exp. Date:



Indications: Recommended for the vaccination of healthy, susceptible cattle as an aid in the prevention of mastitis caused by *Staphylococcus aureus*.

Composition: A lysed culture of highly antigenic polyvalent somatic antigen containing phage types I, II, III, IV and miscellaneous groups of *Staph. aureus*.

Dosage: Using aseptic technique, inject 5 mL intramuscularly. Repeat in 14 days. Follow with a single 5 mL booster dose each 5 to 6 months. Start vaccinating all heifers at 6 months of age.

Precautions: Store out of direct sunlight at a temperature between 35-45°F (2-7°C). Avoid freezing. Shake well before using. Use

Staphylococcus Aureus Bacterin Lysigin®

10 Doses/50mL

entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur.

Antidote: Epinephrine

**Boehringer Ingelheim
Vetmedica, Inc.**
St. Joseph, MO 64506 U.S.A.
U.S. Veterinary License No. 124

Code 104-121; 04101-03



**Boehringer
Ingelheim**

Boehringer Ingelheim Vetmedica, Inc.

2621 North Belt Highway • St. Joseph, MO 64506-2002

Information 800-325-9167 • Customer Service 800-325-9167 • Fax 816-236-2717

www.bi-vetmedica.com

BI 1041-8R-1 12/01