


Serial No.:
Exp. Date:

This product has been shown to be effective for the vaccination of healthy pigs 3 weeks of age or older against *Mycoplasma hyorhinis*. The duration of immunity is at least 7 weeks following vaccination. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

Composition: Prepared from chemically inactivated cultures of *M. hyorhinis*, combined with an adjuvant to stimulate the immune response.

Directions and dosage: Shake well before use. Using aseptic technique, inject a single 2 mL dose intramuscularly. The need for annual booster vaccinations has not been established for this product; consultation with a veterinarian is recommended.

Precautions: Store at 35-46°F (2-8°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 60 days before slaughter. Do not mix with



other products. In case of anaphylactoid reaction, administer epinephrine. The vaccine may induce mild to moderate transient adverse reactions, particularly in young piglets following vaccination. This product has not been tested in pregnant animals. In case of human exposure, contact a physician.

Ingelvac® and MycoMAX™ are trademarks of Boehringer Ingelheim Vetmedica GmbH, used under license.

US 9,273,281 and other patents pending, used under license.

Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, MO 64506
Phone: 866-638-2226
VLN/PCN 124/27H8.00

 **Boehringer
Ingelheim**

127601-00