ToDAY®

Date Fecha	Cow ID Vaca ID	Affected Quarter El Cuarto Affectado				Initials Iniciales	Grade of Mastitis Grado de Mastitis (1, 2, ó 3)
		LH	LF	RH	RF		

Residue Warnings:

1. Milk that has been taken from animals during treatment and for 96 hours after the last treatment must not be used for food.
2. Treated animals must not be slaughtered for food until 4 days after the last treatment.
3. Administration of more than the prescribed dose may lead to residue of antibiotic in milk longer than 96 hours.

Advertencias Sobre Residuos:

1. La leche que se ha extraído de animales en la etapa de tratamiento y durante 96 horas después del último tratamiento no se debe utilizar para alimentos.
2. Los animales tratados no se deben faenar para alimento hasta 4 días después del último tratamiento.
3. La administración de una dosis mayor a la recetada puede dar lugar a residuos de antibióticos en la leche durante más de 96 horas.

Boehringer Ingelheim

ToDAY®

			24 HOUR READ					48 HOUR READ						
Date	Cow ID	Quarter	No Growth	Gram +	Gram -	Contam	Other	No Growth	Gram +	Gram -	Contam	Other	Read By	Dairy Comp

Residue Warnings:

1. Milk that has been taken from animals during treatment and for 96 hours after the last treatment must not be used for food.
2. Treated animals must not be slaughtered for food until 4 days after the last treatment.
3. Administration of more than the prescribed dose may lead to residue of antibiotic in milk longer than 96 hours.

Boehringer Ingelheim